



PROVINCIAL EXAMINATION

JUNE 2023

GRADE 11

MARKING GUIDELINES

MATHEMATICS (PAPER 2)

10 pa

EXEMPLAR

INSTRUCTIONS AND INFORMATION:



- A – ACCURACY
- CA – CONSISTENT ACCURACY
- S – STATEMENT
- R – REASON
- S & R – STATEMENT with REASON

NOTES:

- If a candidate answered a question TWICE, mark only the FIRST attempt.
- If a candidate crossed out an answer and did not redo it, mark the crossed-out answer.
- Consistent accuracy applies to ALL aspects of the marking guidelines.
- It is unacceptable to adopt values/answers in order to solve a problem.

EXEMPLAR

QUESTION 1

1.1	10; 13; 17; 21; 21; 23; 27; 29; 31; 34; 37; 38; 42; 43; 46; 48 Min. = 10 ; Q ₁ = 21 ; Max. = 48 $Q_2 = \frac{29 + 31}{2} = 30 ; Q_3 = \frac{38 + 42}{2} = 40$	✓ Correct Min. Q ₁ and Max. ✓ Correct Q ₂ ✓ Correct Q ₃ (3)
1.2	Grade 11A  Grade 11B 	✓ Correct Min. plot/lower whisker ✓ Correct Max. plot/upper whisker Correct box (3)
1.3	$\bar{x} = \frac{480}{16}$ $\bar{x} = 30$	✓ Fraction ✓ Answer NB: Answer only full marks (2)
1.4	$\sigma = 11,46$	✓✓ Answer (2)
1.5	1 σ interval: (30 - 11,46, 30 + 11,46) (18,54; 41,46) learners achieved over 41,46 marks.	✓ Interval ✓ Answer (2)
1.6	This is a difference in the IQR. IQR = 19 for both classes. However, the middle 50% for Grade 11B was between 21 and 40 compared to only 10 and 30 for Grade 11A. The box for 11B is shifted to the right. Thus, Grade 11B generally performed better.	✓ Comment ✓ IQR value (2)
1.7	75%	✓ Answer (1)
[15]		

QUESTION 2

2.1	<p>Equation of EG:</p> $3x - y + 6 = 0$ $y = 3x + 6$ <p>then $m_{EG} = 3$</p> $\tan \beta = 3$ $\beta = 71,57^\circ$	<ul style="list-style-type: none"> ✓ Gradient ✓ Answer 	(2)
2.2	$x + 3(0) = 8$ $x = 8$ $\therefore (8; 0)$	<ul style="list-style-type: none"> ✓ $x = 0$ ✓ Coordinate form 	(2)
2.3	$M_{DF} = M_{GE}$ diagonals bisect $K\left(\frac{-10+8}{2}; \frac{6+0}{2}\right)$ $K(-1; 3)$ $-1 = \frac{x+1}{2} \quad 3 = \frac{y+9}{2}$ $x = -3 \quad y = -3$ $\therefore E(-3; -3)$	<ul style="list-style-type: none"> ✓ $M_{DE} = M_{GE}$ ✓ Coordinates of K ✓ x-value ✓ y-value 	(4)
2.4	$m_{DF} \times m_{GE}$ $= -\frac{1}{3} \times 3$ $\therefore DF \perp GE$ <p>DGF is a rhombus. (Diagonals bisect perpendicularly)</p>	<ul style="list-style-type: none"> ✓ m_{DF} ✓ $DF \perp GE$ ✓ Reason 	(3)
			[11]

QUESTION 3

3.1	$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ $AC = \sqrt{(1 - (-2))^2 + (5 - (-1))^2}$ $AC = 3\sqrt{5}$	✓ Substitution ✓ Answer	(2)
3.2	$m_{BC} = \frac{5 - 4}{1 - (-3)}$ $m_{BC} = \frac{1}{4}$	✓ Substitution ✓ Answer	(2)
3.3	$m_{BC} = m_{BD} = m_{CD}$ $\frac{1}{4} = \frac{y - 5}{5 - 1}$ or $\frac{1}{4} = \frac{y - 4}{5 - (-3)}$ $y = 6$	✗ Equal gradient ✓ Substitution ✓ $y = 6$	(3)
3.4	$m_{AH} = -4$ (AH \perp BC) $y - (-1) = -4(x - (-2))$ $y = -4x - 9$	✓ $m_{AH} = -4$ ✓ Substitute A ✓ Equation	(3)

[10]

QUESTION 4

4.1	4.1.1	$x^2 + (-24)^2 = 25^2$ $x = -$	✓ Substitute into Pythagoras ✓ Answer	(2)
	4.1.2	(a)	✓ Answer	(1)
		(b)	✓ Reduction ✓ Simplified answer	(2)

		(c)	$\begin{aligned} \tan^2(-\beta) &= \tan^2 \beta \\ &= \left(\frac{-24}{-7}\right)^2 \\ &= \frac{576}{49} \end{aligned}$	<ul style="list-style-type: none"> ✓ Reduction ✓ Simplified answer 	
					(2)
4.1.3		(a)	$\begin{aligned} \sin \beta &= \frac{-24}{25} \\ \frac{y}{15} &= \frac{-24}{25} \\ y &= \frac{-72}{5} \\ \cos \beta &= \frac{-7}{25} \\ \frac{x}{15} &= \frac{-7}{25} \\ x &= \frac{-21}{5} \end{aligned}$	<ul style="list-style-type: none"> ✓ $\sin \beta = \frac{-24}{25}$ ✓ Equate sine ratios ✗ $\cos \beta = \frac{-7}{15}$ ✗ Equate cosine ratios 	
					(4)
		(b)	<p>ANSWER</p> $\begin{aligned} \text{Area}_{\triangle \text{ROT}} &= \frac{1}{2}(12)\left(\frac{72}{5}\right) \\ \text{Area}_{\triangle \text{ROT}} &= 86.4 \text{ units}^2 \end{aligned}$ <p>ALTERNATIVE</p> $\begin{aligned} OT &= 15, \quad OR = 12 \\ A &= 253,75^\circ, \dots^\circ \\ \text{Area}_{\triangle \text{ROT}} &= \frac{1}{2}(15)(12)\sin(106,26^\circ) \\ \text{Area}_{\triangle \text{ROT}} &= 86.4 \text{ units}^2 \end{aligned}$	<ul style="list-style-type: none"> ✓ Formula for area Δ ✓ Base = 12 ✓ Height = $\frac{72}{5}$ ✓ Area ✓ OT length ✓ $106,26^\circ$ ✓ Substitute sine area formula ✓ Area 	
					(4)

4.2	$\begin{aligned} & \frac{\tan 225^\circ + \sin(180^\circ - \theta) \cos(90^\circ + \theta)}{\cos(90^\circ - \theta) \sin(-\theta - 540^\circ)} \\ &= \frac{1 + \sin \theta \cdot -\sin \theta}{\sin \theta \cdot \sin \theta} \\ &= \frac{1 - \sin^2 \theta}{\sin^2 \theta} \\ &= \frac{\cos^2 \theta}{\sin^2 \theta} \\ &= \tan^2 \theta \end{aligned}$	✓ $\tan 225^\circ = 1$ ✓ $\sin(180^\circ - \theta) = \sin \theta$ ✓ $\cos(90^\circ + \theta) = -\sin \theta$ ✓ $\sin(-\theta - 540^\circ) = \sin \theta$ ✓ $1 - \sin^2 \theta = \cos^2 \theta$ ✓ $\frac{\cos^2 \theta}{\sin^2 \theta} = \tan^2 \theta$	(6)
			[21]

QUESTION 5

5.1	$A =$	✓ Answer	(1)
5.2	Period = 120°	✓ Answer	(1)
5.3	$a = -1$ $p = -60^\circ$	✓ Value of a ✓ Value of p	(2)
5.4	$-120^\circ \leq x \leq 60^\circ$ or $x = -240^\circ$ or $x = 240^\circ$	✓ Interval ✓ $-240^\circ; 240^\circ$	(2)
5.5	$p = 30^\circ$	✓✓ Value of p	(2)
			[8]


QUESTION 6

6.1	$\tan\left(\frac{\theta}{2}\right) = \frac{r}{R}$ $R \cdot \tan\left(\frac{\theta}{2}\right) = r$	✓ Substitution ✓ Manipulation	(2)
6.2	$V_{hemisphere} = \frac{1}{2} \cdot \frac{4}{3} \pi R^3$ $V_{hemisphere} = \frac{2}{3} \pi R^3$ $V_{cone} = \frac{1}{3} \pi r^2 R$ $V_{cone} = \frac{1}{3} \pi \left(R \cdot \tan\left(\frac{\theta}{2}\right)\right)^2 R$ $V_{cone} = \frac{2}{3} \pi \cdot R^3 \tan^2\left(\frac{\theta}{2}\right)$ $V_{remaining} = \frac{2}{3} \pi R^3 - \frac{1}{3} \pi \cdot R^3 \tan^2\left(\frac{\theta}{2}\right)$ $V_{remaining} = \frac{\pi R^3}{3} \left(2 - \tan^2\left(\frac{\theta}{2}\right)\right)$	✓ $\frac{1}{2}$ Volume Sphere ✓ Substitution from 6.1 ✓ Subtract volumes (METHOD) ✓ Factors	
6.3	$V_{cone} = \frac{1}{3} \pi \cdot R^3 \tan^2\left(\frac{\theta}{2}\right)$ $V_{cone} = \frac{1}{3} \pi (11)^3 \tan^2\left(\frac{36^\circ}{2}\right)$ $V_{cone} = 147,1 \text{ cm}^3$	✓ $V_{cone} = \frac{1}{3} \pi \cdot R^3 \tan^2\left(\frac{\theta}{2}\right)$ ✓ Substitute $R = 11$ and $\theta = 36^\circ$ ✓ Volume of wax	(3)


QUESTION 7

7.1	7.1.1 $\hat{A} = 40^\circ$	tan chord theorem	✓ S ✓ R	(2)
	7.1.2 $K_1 = 90^\circ$	\angle 's in a semi circle	✓ S ✓ R	(2)
	7.1.3 $\hat{P}_1 = 25^\circ$	ext \angle of a Δ	✓ S ✓ R	(2)
7.2	$\hat{T}_1 = 65^\circ$ TC = QC	ext \angle of Δ sides opp equal \angle 's	✓ R ✓ R	(2)
				[8]

QUESTION 8



8.1	<p>Construct UO and extend.</p> <p>In $\triangle TOU$ $\hat{O}_1 = \hat{T} + \hat{U}_1$ $\hat{T} = \hat{U}_1$ $\therefore \hat{O}_1 = \hat{U}_1$</p> <p>exterior \angle of \triangle \angle's of equ radii</p> <p>Similarly in $\triangle TOU$ $\therefore \hat{O}_2 = \hat{U}_2$ $\therefore \hat{O}_1 + \hat{O}_2 = 2\hat{U}_1$ $\hat{O}_1 + \hat{O}_2 = 2(\hat{U}_1 + \hat{U}_2)$ $TOU = TVU$</p>	<p>✓ Construction</p> <p>✓ SR</p> <p>✓ SR</p> <p>✓ S</p> <p>✓ Conclusion</p>	(5)



8.2	8.2.1		Two pairs of adjacent sides e	✓ R	(1)
	8.2.2	(a)	\angle at centre = $2 \times \angle$ at circumference	✓ R	(1)
		(b)	ext \angle of cyclic quad	✓ R	(1)
		(c)	corr \angle 's, $OB \parallel DM$	✓ R	(1)
		(d)	opp \angle 's, rhombus	✓ R	
			ALTERNATIVE		
			a) \angle 's, $OB \parallel DM$		(1)
	8.2.3		$D_2 = 90^\circ$ $\hat{D}_2 = 90^\circ - x$ $\hat{x} = 90^\circ$ $\hat{C} = 90^\circ - x$ $\therefore \hat{D}_2 = \hat{C}$	$\tan \perp \text{rad}$ $\tan \perp \text{rad}$ $\text{int } \angle \text{s of } \Delta$ both $90^\circ - x$	✓ R ✓ S ✓ SR ✓ Conclusion (4)
	8.2.4		$\hat{F}_2 = 90^\circ$ $AF = FD$ $AB = BC$	corr \angle s, $FB \parallel DM$ line from centre \perp chord line through midpt \parallel to 2 nd side	✓ SR ✓ S ✓ R ✓ R (4)
					[18]
					TOTAL: 100